



## References

Layne JE, Jepson LH, Carite AM, Parkin CG, Bergenstal RM. Long-Term Improvements in Glycemic Control with Dexcom CGM Use in Adults with Noninsulin-Treated Type 2 Diabetes. *Diabetes Technol Ther*. 2024 Jun 21. doi: 10.1089/dia.2024.0197. Epub ahead of print.

Norman GJ, Fernandes J, Nemlekar P, Andrade SB, Lupton L, Berk A. Initiating continuous glucose monitoring is associated with improvements in glycemic control and reduced health care resource utilization for people with diabetes in a large US-insured population: A real-world evidence study. *J Manag Care Spec Pharm*. 2024:1-10.

Grace T, Layne J, Hicks C, Green C, Walker, T. The Dexcom Community Glucose Monitoring Project for People with T2D: 1-Year Outcomes [abstract]. Presented at: ADA 84th Scientific Sessions; Orlando, FL; June 21–24, 2024. Abstract 982-P.

Alshannaq H, Norman G, Lynch P. Cost-Effectiveness of Real-time CGM vs. Self-Monitoring of Blood Glucose in People with Insulin-treated Type 2 Diabetes from a US Payor Perspective [abstract]. Presented at: ADA 84th Scientific Sessions; Orlando, FL; June 21–24, 2024. Abstract 982-P. Abstract 1035-P.

Garg SK, Hirsch IB, Repetto E, Snell-Bergeon J, Ulmer B, Perkins C, Bergenstal RM. Impact of continuous glucose monitoring on hospitalizations and glucose control in people with type 2 diabetes: real-world analysis. *Diabetes Obes Metab*. 2024 Sep 12. doi: 10.1111/dom.15866. Epub ahead of print.



## Acronyms

CGM=continuous glucose monitoring

T1D=type 1 diabetes

T2D=type 2 diabetes

NIT=non-insulin therapy

NIIT=non-intensive insulin therapy

IIT=intensive insulin therapy

GMI=glucose management indicator

TIR=time in range

TITR=time in tight range

HEDIS=Healthcare Effectiveness Data and Information Set

ADA=American Diabetes Association

rtCGM=real-time continuous glucose monitoring

NNT=number needed to treat

ACH=all-cause hospitalization

ADH=acute diabetes-related event requiring hospitalization

ADER=acute diabetes-related event requiring emergency room visit